

Parameterized algorithms and data reduction for safe convoy routing

René van Bevern

Department of Mechanics and Mathematics, Novosibirsk State University
Ulitsa Pirogova 2, 630090 Novosibirsk, Russian Federation

Sobolev Institute of Mathematics of the Siberian Branch of the Russian Academy of Sciences
Prospekt Akademika Koptyuga 4, 630090 Novosibirsk, Russian Federation

rvb@nsu.ru

 <https://orcid.org/0000-0002-4805-218X>

Till Fluschnik

Institut für Softwaretechnik und Theoretische Informatik, TU Berlin
Ernst-Reuter-Platz 7, 10587 Berlin, Germany

till.fluschnik@tu-berlin.de

Oxana Yu. Tsidulko

Sobolev Institute of Mathematics of the Siberian Branch of the Russian Academy of Sciences
Prospekt Akademika Koptyuga 4, 630090 Novosibirsk, Russian Federation

Department of Mechanics and Mathematics, Novosibirsk State University
Ulitsa Pirogova 2, 630090 Novosibirsk, Russian Federation

tsidulko@math.nsc.ru

Abstract

We study a problem that models safely routing a convoy through a transportation network, where any vertex adjacent to the travel path of the convoy requires additional precaution: Given a graph $G = (V, E)$, two vertices $s, t \in V$, and two integers k, ℓ , we search for a simple s - t -path with at most k vertices and at most ℓ neighbors. We study the problem in two types of transportation networks: graphs with small crossing number, as formed by road networks, and tree-like graphs, as formed by waterways. For graphs with constant crossing number, we provide a subexponential $2^{O(\sqrt{n})}$ -time algorithm and prove a matching lower bound. We also show a polynomial-time data reduction algorithm that reduces any problem instance to an equivalent instance (a so-called problem kernel) of size polynomial in the vertex cover number of the input graph. In contrast, we show that the problem in general graphs is hard to preprocess. Regarding tree-like graphs, we obtain a $2^{O(\text{tw})} \cdot \ell^2 \cdot n$ -time algorithm for graphs of treewidth tw , show that there is no problem kernel with size polynomial in tw , yet show a problem kernel with size polynomial in the feedback edge number of the input graph.

2012 ACM Subject Classification Mathematics of computing → Discrete mathematics → Graph theory → Graph algorithms

Keywords and phrases NP-hard problem · fixed-parameter tractability · problem kernelization · shortest path · secluded solution

Digital Object Identifier [10.4230/OASICS.ATMOS.2018.10](https://doi.org/10.4230/OASICS.ATMOS.2018.10)

Related Version Full version available at [arXiv:1806.09540](https://arxiv.org/abs/1806.09540)

Funding Till Fluschnik was supported by the German Research Foundation, grant NI 369/18. René van Bevern and Oxana Yu. Tsidulko were supported by the Russian Foundation for Basic Research (RFBR), grant 18-501-12031 NNIO_a, while working on Sections 2.2, 2.3, 3.3, and 3.2.



© René van Bevern, Till Fluschnik, Oxana Yu. Tsidulko;
licensed under Creative Commons License CC-BY

18th Workshop on Algorithmic Approaches for Transportation Modelling, Optimization, and Systems (ATMOS 2018).

Editors: Ralf Borndörfer and Sabine Storandt; Article No. 10; pp. 10:1–10:19

[OpenAccess Series in Informatics](#)



OASICS Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

While working on Sections 2.1 and 3.1, René van Bevern was supported by RFBR grant 16-31-60007 mol_a_dk and Oxana Yu. Tsidulko was supported by the Ministry of Science and Higher Education of the Russian Federation under the 5-100 Excellence Programme.

1 Introduction

Finding shortest paths is a fundamental problem in route planning and has extensively been studied with respect to efficient algorithms, including data reduction and preprocessing [1]. In this work, we study the following NP-hard variant of finding shortest s - t -paths.

► **Problem 1.1** (SHORT SECLUDED PATH (SSP)).

Input: An undirected, simple graph $G = (V, E)$ with two distinct vertices $s, t \in V$, and two integers $k \geq 2$ and $\ell \geq 0$.

Question: Is there an s - t -path P in G such that $|V(P)| \leq k$ and $|N(V(P))| \leq \ell$?

Herein, $V(P)$ denotes the set of vertices on path P and $N(V(P))$ denotes their set of neighbors (not lying on P).

The problem can be understood as finding short and safe routes for a convoy through a transportation network: each neighbor of the convoy’s travel path requires additional precaution. Thus, we seek to minimize not only the length of the convoy’s travel path, but also its number of neighbors. In our work, we study the above basic, unweighted variant, as well as a weighted variant of the problem, in which each vertex has two weights: one counts towards the path length, the other models the cost of precaution that has to be taken when the vertex occurs as the neighbor of the travel path.

Almost planar and tree-like transportation networks. The focus of our work is two-fold. Firstly, since the problem is NP-hard, we search for efficient algorithms in graphs that are likely to occur as transportation networks: almost planar graphs, which occur as road networks, and tree-like graphs, which arise as waterways (ignoring the few man-made canals, natural river networks form forests [23]). Secondly, given the effect that preprocessing and data reduction had to fundamental routing problems like finding shortest paths [1], we study the possibilities of polynomial-time data reduction with *provable performance guarantees* for SSP.

In order to measure the running time of our algorithms with respect to the “degree of planarity” or the “tree-likeness” of a graph, as well as to analyze the power of data reduction algorithms, we employ parameterized complexity theory, which provides us with the concepts of *fixed-parameter algorithms* and *problem kernelization* [12, 16, 20, 39]. Fixed-parameter algorithms have recently been applied to numerous NP-hard routing problems [2, 3, 5, 15, 26–30, 41, 42]. In particular, they led to subexponential-time algorithms for fundamental NP-hard routing problems in planar graphs [33] and to algorithms for hard routing problems that work efficiently on real-world data [3].

Fixed-parameter algorithms. The main idea of fixed-parameter algorithms is to accept the exponential running time seemingly inherent to solving NP-hard problems, yet to restrict the combinatorial explosion to a parameter of the problem, which can be small in applications. We call a problem *fixed-parameter tractable* if it can be solved in $f(k) \cdot n^{O(1)}$ time on inputs of length n and some function f depending only on some parameter k . In contrast to an algorithm that merely runs in polynomial time for fixed k , fixed-parameter algorithms can solve NP-hard problems quickly if k is small.

■ **Table 1** Overview of our results. Herein, n , tw , vc , fes , cr , and Δ denote the number of vertices, treewidth, vertex cover number, feedback edge number, the crossing number, and maximum degree of the input graph, respectively. “const.” abbreviates “constant”.

	On almost planar graphs (Sec. 2)	on tree-like graphs (Sec. 3)
exact solution	$2^{O(\sqrt{n})}$ time in graphs with const. cr (Thm. 2.1)	$2^{O(tw)} \cdot \ell^2 \cdot n$ time (Thm. 3.2)
problem kernel	size $vc^{O(r)}$ in $K_{r,r}$ -free graphs (Thm. 2.5)	size $fes^{O(1)}$ (Thm. 3.13)
lower bounds	No kernel with size $\text{poly}(vc + r)$ in $K_{r,r}$ -free graphs and WK[1]-hard when parameterized by $vc + r$ (Thm. 2.14)	No kernel with size $\text{poly}(tw + k + \ell)$ even in planar graphs with const. Δ (Thm. 3.10)

Provably effective polynomial-time data reduction. Kernelization allows for provably effective polynomial-time data reduction. Note that a result of the form “our polynomial-time data reduction algorithm reduces the input size by at least one bit, preserving optimality of solutions” is impossible for NP-hard problems unless $P = NP$. In contrast, a kernelization algorithm reduces a problem instance into an equivalent one (the *problem kernel*) whose size depends only (ideally polynomially) on some problem parameter. Problem kernelization has been successfully applied to obtain effective polynomial-time data reduction algorithms for many NP-hard problems [25, 34] and also led to techniques for proving the limits of polynomial-time data reduction [6, 7, 38].

1.1 Our contributions

We study SSP (and a weighted variant) in two main classes of graphs: almost planar graphs and tree-like graphs. We refer to Table 1 for an overview on our main results. Regarding almost planar graphs, in graphs of constant crossing number, we show that (even the weighted version of) SSP is solvable in subexponential $2^{O(\sqrt{n})}$ -time. Moreover, we prove that SSP is not solvable in $2^{o(\sqrt{n})}$ -time in planar graphs unless the Exponential Time Hypothesis fails. In $K_{r,r}$ -free graphs, which comprise the graphs with crossing number $O(r^3)$ [40], we show a problem kernel for SSP with size $vc^{O(r)}$, where vc is the vertex cover number of the input graph. We prove that, unless the polynomial-time hierarchy collapses, there is no problem kernel of size polynomial in $vc + r$. Moreover, we prove that, unless the classes FPT and WK[1] coincide, SSP does not even allow for *Turing kernels* with size polynomial in $vc + r$; that is, we could not solve SSP in polynomial time even if we precomputed all answers to subproblems of size polynomial in $vc + r$ and could look them up in constant time. Regarding tree-like graphs, we prove that SSP is solvable in $2^{O(tw)} \cdot \ell^2 \cdot n$ time in graphs of treewidth tw and that there is no problem kernel with size polynomial in tw . Instead, we show a problem kernel of size $fes^{O(1)}$, where fes is the feedback edge number of the input graph.

Due to space constraints, results marked with (\star) are deferred to a full version of the paper.

1.2 Related work

Several classical graph optimization problems have been studied in the “secluded” (small closed neighborhood) and the “small secluded” (small set with small open neighborhood) variants [4]. Luckow and Fluschnik [37] first defined SSP and analyzed its parameterized complexity with respect to the parameters k and ℓ . In contrast, we study problem parameters that describe the structure of the input graphs and are small in transportation networks. Chechik et al. [11] introduced the SECLUDED PATH problem, that, given an undirected

graph $G = (V, E)$ with two designated vertices $s, t \in V$, vertex-weights $w : V \rightarrow \mathbb{N}$, and two integers $k, C \in \mathbb{N}$, asks whether there is an s - t -path P such that the size of the *closed* neighborhood $|N[V(P)]| \leq k$ and the weight of the closed neighborhood $w(N[V(P)]) \leq C$. Fomin et al. [21], in particular, prove that SECLUDED PATH does not admit problem kernels with size polynomial in the vertex cover number vc . Our negative results on kernelization for SSP are significantly stronger: not only do we show that there is no problem kernel of size polynomial in $vc + r$ even in bipartite $K_{r,r}$ -free graphs, we also show that SSP is WK[1]-hard parameterized by $vc + r$. Golovach et al. [24] studied the “small secluded” scenario for finding connected induced subgraphs parameterized by the size ℓ of the open neighborhood. Their results obviously does not generalize to SSP, since SSP is NP-hard even for $\ell = 0$ [37].

1.3 Preliminaries

Graph Theory. We use basic notation from graph theory [14]. We study simple, finite, undirected graphs $G = (V, E)$. We denote by $V(G) := V$ the set of *vertices* of G and by $E(G) := E$ the set of *edges* of G . We denote $n := |V|$ and $m := |E|$. For any subset $U \subseteq V$ of vertices, we denote by $N_G(U) = \{w \in V \setminus U \mid \exists v \in U : \{v, w\} \in E\}$ the *open neighborhood* of U in G . When the graph G is clear from the context, we drop the subscript G . A set $U \subseteq V$ of vertices is a *vertex cover* if every edge in E has an endpoint in U . The size of a minimum vertex cover is called *vertex cover number* $vc(G)$ of G . A set $F \subseteq E$ of edges is a *feedback edge set* if the graph $(G, E \setminus F)$ is a forest. The minimum size of a feedback edge set in a connected graph is $m - n + 1$ and is called the *feedback edge number* $fes(G)$ of G . The *crossing number* $cr(G)$ of G is the minimum number of crossings in any drawing of G into the plane (where only two edges are allowed to cross in each point). A path $P = (V, E)$ is a graph with vertex set $V = \{x_0, x_1, \dots, x_p\}$ and edge set $E = \{\{x_i, x_{i+1}\} \mid 0 \leq i < p\}$. We say that P is an x_0 - x_p -path of length p . We also refer to x_0, x_p as the *end points* of P , and to all vertices $V \setminus \{x_0, x_p\}$ as the *inner* vertices of P . A $K_{r,r}$ is a complete bipartite graph $G = (U \uplus V, E)$ with $|U| = |V| = r$. We say that a graph is $K_{r,r}$ -free if it does not contain $K_{r,r}$ as a subgraph.

Parameterized Complexity Theory. For more details on parameterized complexity, we refer to the text books [12, 16, 20, 39]. Let Σ be a finite alphabet. A *parameterized problem* L is a subset $L \subseteq \Sigma^* \times \mathbb{N}$. An instance $(x, k) \in \Sigma^* \times \mathbb{N}$ is a *yes-instance* for L if and only if $(x, k) \in L$. We call x the *input* and k the *parameter*.

► **Definition 1.2** (fixed-parameter tractability, FPT). A parameterized problem $L \subseteq \Sigma^* \times \mathbb{N}$ is *fixed-parameter tractable* if there is a *fixed-parameter algorithm* deciding $(x, k) \in L$ in time $f(k) \cdot |x|^{O(1)}$. The complexity class *FPT* consists of all fixed-parameter tractable problems.

► **Definition 1.3** (kernelization). Let $L \subseteq \Sigma^* \times \mathbb{N}$ be a parameterized problem. A *kernelization* is an algorithm that maps any instance $(x, k) \in \Sigma^* \times \mathbb{N}$ to an instance $(x', k') \in \Sigma^* \times \mathbb{N}$ in $\text{poly}(|x| + k)$ time such that

- (i) $(x, k) \in L \iff (x', k') \in L'$, and
- (ii) $|x'| + k' \leq f(k)$ for some computable function f .

We call (x', k') the *problem kernel* and f its *size*.

Basic observations. We may assume our input graph to be connected due to the following obviously correct and linear-time executable data reduction rule.

► **Reduction Rule 1.4.** If G has more than one connected component, then delete all but the component containing both s and t or return no if such a component does not exist.

2 Almost planar graphs

Many transportation networks such as rail and street networks are planar or at least have a small *crossing number*—the minimum number of edge crossings in a plane drawing of a graph. Unfortunately, SSP remains NP-hard even in planar graphs with maximum degree four and $\ell = 0$ [37].

In this section, we present algorithms for SSP in graphs with constant crossing number. These, in particular, apply to planar graphs. First, in Section 2.1, we present a subexponential-time algorithm and a matching lower bound. Second, in Section 2.2, we present a provably effective data reduction algorithm. Finally, in Section 2.3, we show the limits of data reduction algorithms for SSP in graphs with small but non-constant crossing number.

2.1 A subexponential-time algorithm

In this section, we describe how to solve SSP in subexponential time in graphs with constant crossing number.

► **Theorem 2.1.** *SHORT SECLUDED PATH is solvable in $2^{O(\sqrt{n})}$ time on graphs with constant crossing number.*

We will also see a matching lower bound. To prove Theorem 2.1, we exploit that graphs with constant crossing number are *H-minor free* for some graph *H*.

► **Definition 2.2** (graph minor). A graph *H* is a *minor* of a graph *G* if *H* can be obtained from *G* by a sequence of vertex deletions, edge deletions, and edge contractions. If a graph *G* does not contain *H* as a minor, then *G* is said to be *H-minor free*.

Bokal et al. [10] showed that, if a graph *G* contains $K_{r,r}$ as a minor, then the crossing number of *G* is $\text{cr}(G) \geq \frac{1}{2}(r-2)^2$. Thus, any graph *G* is $K_{r,r}$ -minor free for $r > \sqrt{2\text{cr}(G)} + 2$, which goes in line with the well-known fact that planar graphs are $K_{3,3}$ -minor free [43]. Demaine and Hajiaghayi [13] showed that, for any graph *H*, all *H*-minor free graphs have treewidth $\text{tw} \in O(\sqrt{n})$.¹ To prove Theorem 2.1, it thus remains to show that SSP is solvable in $2^{O(\text{tw})} \cdot \text{poly}(n)$ time, which is the main technical work deferred to Section 3.1.

Complementing Theorem 2.1, we can show a matching lower bound using the Exponential Time Hypothesis (ETH).

► **Conjecture 2.3** (Exponential Time Hypothesis (ETH), Impagliazzo et al. [32]). *There is a constant c such that n -variable 3-SAT cannot be solved in $2^{c(n+m)}$ time.*

The ETH was introduced by Impagliazzo et al. [32] and since then has been used to prove running time lower bounds for various NP-hard problems (we refer to Cygan et al. [12, Chapter 14] for an overview). We use it to prove that Theorem 2.1 can be neither significantly improved in planar graphs nor generalized to general graphs.

► **Theorem 2.4.** *Unless the Exponential Time Hypothesis fails, SHORT SECLUDED PATH has no $2^{o(\sqrt{n})}$ -time algorithm in planar graphs and no $2^{o(n+m)}$ -time algorithm in general.*

Proof. Assume that there is a $2^{o(\sqrt{n})}$ -time algorithm for SSP in planar graphs and a $2^{o(n)}$ -time algorithm for SSP in general graphs. Luckow and Fluschnik [37] give a polynomial-time

¹ In fact, they showed $\text{tw} \in O(\sqrt{q})$ for any graph parameter q that is $\Omega(p)$ on a $(\sqrt{p} \times \sqrt{p})$ -grid and does not increase when taking minors. For example, the vertex cover number or feedback vertex number.

many-one reduction from HAMILTONIAN CYCLE to SSP that maintains planarity and increases the number of vertices and edges by at most a constant. Thus, we get a $2^{o(\sqrt{n})}$ -time algorithm for HAMILTONIAN CYCLE in planar graphs and a $2^{o(n+m)}$ -time algorithm in general graphs. This contradicts ETH [12, Theorems 14.6 and 14.9]. ◀

2.2 Effective data reduction

In the previous section, we have shown a subexponential-time algorithm for SSP in graphs with constant crossing number. There, we exploited the fact that graphs with crossing number cr are $K_{r,r}$ -minor free for $r > \sqrt{2cr} + 2$. Of course, this means that they neither contain $K_{r,r}$ as subgraph (indeed, one can show this even for $r \geq 3.145 \cdot \sqrt[3]{cr}$ using bounds from Pach et al. [40]).

In this section, we show how to reduce any instance of SSP in $K_{r,r}$ -free graphs to an equivalent instance with size polynomial in the vertex cover number of the input graph. In the next section, we prove that this does not generalize to general graphs.

► **Theorem 2.5.** *For each constant $r \in \mathbb{N}$, SHORT SECLUDED PATH in $K_{r,r}$ -free graphs admits a problem kernel with size polynomial in the vertex cover number of the input graph.*

The proof of Theorem 2.5 consists of three steps. First, in linear time, we transform an n -vertex instance of SSP into an equivalent instance of an auxiliary vertex-weighted version of SSP with $O(vc^r)$ vertices. Second, using a theorem of Frank and Tardos [22], in polynomial time, we reduce the vertex weights to $2^{O(vc^{3r})}$ so that the total instances size (in bits) becomes $O(vc^{4r})$. Finally, since SSP is NP-complete in planar, and, hence, in $K_{3,3}$ -free graphs, we can, in polynomial time, reduce the shrunk instance back to an instance of the unweighted SSP in $K_{r,r}$ -free graphs. Due to the polynomial running time of the reduction, there is at most a polynomial blow-up of the instance size.

Our auxiliary variant of SSP allows each vertex to have two weights: one weight counts towards the length of the path, the other counts towards the number of neighbors:

► **Problem 2.6** (VERTEX-WEIGHTED SHORT SECLUDED PATH (VW-SSP)).

Input: An undirected, simple graph $G = (V, E)$ with two distinct vertices $s, t \in V$, two integers $k \geq 2$ and $\ell \geq 0$, and vertex weights $\kappa : V \rightarrow \mathbb{N}$ and $\lambda : V \rightarrow \mathbb{N}$.

Question: Does G have an s - t -path P with $\sum_{v \in V(P)} \kappa(v) \leq k$ and $\sum_{v \in N(V(P))} \lambda(v) \leq \ell$?

Note that an instance of SSP can be considered to be an instance of VW-SSP with unit weight functions κ and λ . Our data reduction will be based on removing *twins*.

► **Definition 2.7** (twins). Two vertices u and v are called (*false*) *twins* if $N(u) = N(v)$.

As the first step towards proving Theorem 2.5, we will show that the following data reduction rule, when applied to a $K_{r,r}$ -free instance of SSP for constant r , leaves us with an instance of VW-SSP with $O(vc^r)$ vertices.

► **Reduction Rule 2.8.** Let $(G, s, t, k, \ell, \kappa, \lambda)$ be an VW-SSP instance with unit weights, where $G = (V, E)$ is a $K_{r,r}$ -free graph.

For each maximal set $U \subseteq V \setminus \{s, t\}$ of twins such that $|U| > r$, delete $|U| - r + 1$ vertices of U from G , and, for an arbitrary remaining vertex $v \in U$, set $\lambda(v) := |U| - r$ and $\kappa(v) := k + 1$.

► **Lemma 2.9** (*). *Reduction Rule 2.8 is correct and can be applied in linear time.*

We now prove a size bound for the instances remaining after Reduction Rule 2.8.

► **Proposition 2.10.** *Applied to an instance of SSP with a $K_{r,r}$ -free graph with vertex cover number vc , Reduction Rules 2.8 and 1.4 yield an instance of VW-SSP on at most $(vc + 2) + r(vc + 2)^r$ vertices in linear time.*

Proof. Let $(G', s, t, k, \ell, \lambda', \kappa')$ be the instance obtained from applying Reduction Rules 2.8 and 1.4 to an instance $(G, s, t, k, \ell, \lambda, \kappa)$.

Let C be a minimum-cardinality vertex cover for G' that contains s and t , and let the vertex set of G' be $V = C \uplus Y$. Since G' is a subgraph of G , one has $|C| \leq vc(G') + 2 \leq vc(G) + 2 = vc + 2$. It remains to bound $|Y|$. To this end, we bound the number of vertices of degree at least r in Y and the number of vertices of degree exactly i in Y for each $i \in \{0, \dots, r-1\}$. Note that vertices in Y have neighbors only in C .

Since Reduction Rule 1.4 has been applied, there are no vertices of degree zero in Y .

Since Reduction Rule 2.8 has been applied, for each $i \in \{1, \dots, r-1\}$ and each subset $C' \subseteq C$ with $|C'| = i$, we find at most r vertices in Y whose neighborhood is C' . Thus, for each $i \in \{1, \dots, r-1\}$, the number of vertices with degree i in Y is at most $r \cdot \binom{|C|}{i}$.

Finally, since G is $K_{r,r}$ -free, any r -sized subset of the vertex cover C has at most $r-1$ common neighbors. Hence, since vertices in Y have neighbors only in C , the number of vertices in Y of degree greater or equal to r is at most $(r-1) \cdot \binom{|C|}{r}$. We conclude that

$$|V'| \leq |C| + (r-1) \cdot \binom{|C|}{r} + r \cdot \sum_{i=1}^{r-1} \binom{|C|}{i} \leq (vc + 2) + r(vc + 2)^r. \quad \blacktriangleleft$$

This completes the first step of the proof of Theorem 2.5. Note that our data reduction works by “hiding” an unbounded number of twins in vertices of unbounded weights. The second step is thus reducing the weights of an VW-SSP instance. To this end, we are going to apply a theorem by Frank and Tardos [22], which was successfully applied in kernelizing weighted problems before [17].

► **Proposition 2.11** (Frank and Tardos [22]). *There is an algorithm that, on input $w \in \mathbb{Q}^d$ and integer N , computes in polynomial time a vector $\bar{w} \in \mathbb{Z}^d$ with $\|\bar{w}\|_\infty \leq 2^{4d^3} N^{d(d+2)}$ such that $\text{sign}(w^\top b) = \text{sign}(\bar{w}^\top b)$ for all $b \in \mathbb{Z}^d$ with $\|b\|_1 \leq N-1$, where*

$$\text{sign}(x) = \begin{cases} +1 & \text{if } x > 0, \\ 0 & \text{if } x = 0, \text{ and} \\ -1 & \text{if } x < 0. \end{cases}$$

► **Observation 2.12.** *For $N \geq 2$, Proposition 2.11 gives $\text{sign}(w^\top e_i) = \text{sign}(\bar{w}^\top e_i)$ for each $i \in \{1, \dots, d\}$, where $e_i \in \mathbb{Z}^d$ is the vector that has 1 in the i -th coordinate and zeroes in the others. Thus, one has $\text{sign}(w_i) = \text{sign}(\bar{w}_i)$ for each $i \in \{1, \dots, d\}$. That is, when reducing a weight vector from w to \bar{w} , Proposition 2.11 maintains the signs of weights.*

We apply Proposition 2.11 and Observation 2.12 to the weights of VW-SSP.

► **Lemma 2.13.** *An instance $I = (G, s, t, k, \ell, \lambda, \kappa)$ of VW-SSP on an n -vertex graph $G = (V, E)$ can be reduced in polynomial time to an instance $I' = (G, s, t, k', \ell', \lambda', \kappa')$ of VW-SSP such that*

- i) $\{k', \kappa'(v), \ell', \lambda'(v)\} \subseteq \{0, \dots, 2^{4(n+1)^3} \cdot (n+2)^{(n+1)(n+3)}\}$, for each vertex $v \in V$, and
- ii) I is a yes-instance if and only if I' is a yes-instance.

Proof. In this proof, we will conveniently denote the weight functions $\lambda, \lambda', \kappa$, and κ' as vectors in \mathbb{N}^n such that $\lambda_i = \lambda(i)$ for each $i \in V$, and similarly for the other weight functions.

We apply Proposition 2.11 with $d = n + 1$ and $N = n + 2$ to the vectors $(\lambda, \ell) \in \mathbb{N}^{n+1}$ and $(\kappa, k) \in \mathbb{N}^{n+1}$ to obtain vectors $(\kappa', k') \in \mathbb{Z}^{n+1}$ and $(\lambda', \ell') \in \mathbb{Z}^{n+1}$ in polynomial time.

(i) This follows from Proposition 2.11 with $d = n + 1$ and $N = n + 2$, and from Observation 2.12 since (λ, ℓ) and (κ, k) are vectors of nonnegative numbers.

(ii) Consider an arbitrary s - t -path P in G and two associated vectors $x, y \in \mathbb{Z}^n$, where

$$x_v = \begin{cases} 1 & \text{if } v \in N(V(P)), \\ 0 & \text{otherwise,} \end{cases} \quad y_v = \begin{cases} 1 & \text{if } v \in V(P) \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

Observe that $\|(x, -1)\|_1 \leq n + 1$ and $\|(y, -1)\|_1 \leq n + 1$. Since $n + 1 \leq N - 1$, Proposition 2.11 gives $\text{sign}((\lambda, \ell)^\top(x, -1)) = \text{sign}((\lambda', \ell')^\top(x, -1))$ and $\text{sign}((\kappa, k)^\top(y, -1)) = \text{sign}((\kappa', k')^\top(y, -1))$, which is equivalent to

$$\sum_{v \in N(V(P))} \lambda(v) \leq \ell \iff \sum_{v \in N(V(P))} \lambda'(v) \leq \ell' \quad \text{and} \quad \sum_{v \in P} \kappa(v) \leq k \iff \sum_{v \in P} \kappa'(v) \leq k'. \quad \blacktriangleleft$$

We have finished two steps towards the proof of Theorem 2.5: we reduced SSP in $K_{r,r}$ -free graphs for constant r to instances of VW-SSP with $O(vc^r)$ vertices using Proposition 2.10 and shrunk its weights to encoding-length $O(vc^{3r})$ using Lemma 2.13. To finish the proof of Theorem 2.5, it remains to reduce VW-SSP back to SSP on $K_{r,r}$ -free graphs.

2.3 Limits of data reduction

In Section 2.2, we have seen that SSP allows for problem kernels with size polynomial in vc if the input graph is $K_{r,r}$ -free for some constant r . A natural question is whether one can loosen the requirement of r being *constant*.

The following Theorem 2.14(i) shows that, under reasonable complexity-theoretic assumptions, this is not the case: we cannot get problem kernels whose size bound depends polynomially on both vc and r . Moreover, the following Theorem 2.14(ii) shows that, unless $\text{WK}[1] = \text{FPT}$, SSP does not even have Turing kernels with size polynomial in $vc + r$ [31]. That is, we could not even solve SSP in polynomial time if we had precomputed all answers to SSP instances with size polynomial in $vc + r$ and could look them up in constant time.

Both results come surprisingly: finding a standard shortest s - t -path is easy, whereas finding a short secluded path in general graphs is so hard that not even preprocessing helps.

► **Theorem 2.14** (\star). *Even in bipartite graphs, SHORT SECLUDED PATH*

i) *has no problem kernel with size polynomial in $vc + r$ unless $\text{coNP} \subseteq \text{NP/poly}$ and*

ii) *is WK[1]-hard when parameterized by $vc + r$,*

where vc is the vertex cover number of the input graph and r is the smallest number such that the input graph is $K_{r,r}$ -free.

The proof exploits that MULTICOLORED CLIQUE is WK[1]-hard parameterized by $k \log n$ [31]:


► **Problem 2.15** (MULTICOLORED CLIQUE).

Input: A k -partite n -vertex graph $G = (V, E)$, where $V = \bigsqcup_{i=1}^k V_i$ for independent sets V_i .

Question: Does G contain a clique of size k ?

We transfer the WK[1]-hardness of MULTICOLORED CLIQUE to SSP using the following type of reduction [31]:

► **Definition 2.16** (polynomial parameter transformation). Let $L, L' \subseteq \Sigma^* \times \mathbb{N}$ be two parameterized problems. A *polynomial parameter transformation* from L to L' is an algorithm that maps any instance $(x, k) \in \Sigma^* \times \mathbb{N}$ to an instance $(x', k') \in \Sigma^* \times \mathbb{N}$ in $\text{poly}(|x| + k)$ time such that



■ **Figure 2.1** Illustration of the polynomial parameter transformation. Non-black (green) vertices indicate the vertices in the vertex cover.

- (i) $(x, k) \in L \iff (x', k') \in L'$, and
- (ii) $k' \leq \text{poly}(k)$.

Our polynomial parameter transformation of MULTICOLORED CLIQUE into SSP uses the following gadget.

► **Definition 2.17** (*z*-binary gadget). A *z*-binary gadget for some power *z* of two is a set $B = \{u_1, u_2, \dots, u_{(2 \log z)}\}$ of vertices. We say that a vertex *v* is *p*-connected to *B* for some $p \in \{0, \dots, z-1\}$ if *v* is adjacent to $u_q \in B$ if and only if there is a “1” in position *q* of the string that consists of the binary encoding of *p* followed by its complement.

► **Example 2.18.** The binary encoding of 5 followed by its complement is 101010. Thus, a vertex *v* is 5-connected to an 8-binary gadget $\{u_1, \dots, u_6\}$ if and only if *v* is adjacent to u_1, u_3 , and u_5 . Also observe that, if a vertex *v* is *q*-connected to a *z*-binary gadget *B*, then *v* is adjacent to exactly half of the vertices of *B*, that is, to $\log z$ vertices of *B*.

The following reduction from MULTICOLORED CLIQUE to SSP is illustrated in Figure 2.1.

► **Construction 2.19.** Let $G = (V, E)$ be a MULTICOLORED CLIQUE instance, where $|V| = n$ and $V = V_1 \uplus V_2 \uplus \dots \uplus V_k$. Without loss of generality, assume that $V_i = \{v_i^1, v_i^2, \dots, v_i^{\tilde{n}}\}$ for each $i \in \{1, \dots, k\}$, where \tilde{n} is some power of two (we can guarantee this by adding isolated vertices to *G*). We construct an equivalent instance (G', s, t, k', ℓ') of SSP, where

$$k' := \binom{k}{2} + 1, \quad \ell' := |E| - \binom{k}{2} + k \log \tilde{n},$$

and the graph $G' = (V', E')$ is as follows. The vertex set V' consists of vertices *s*, *t*, a vertex v_e for each edge $e \in E$, vertices w_h for $h \in \{1, \dots, \binom{k}{2} - 1\}$, and mutually disjoint \tilde{n} -binary vertex gadgets B_1, \dots, B_k , each vertex in which has $\ell' + 1$ neighbors of degree one. We denote

$$\begin{aligned} E^* &:= \{v_e \in V' \mid e \in E\} & B &:= B_1 \uplus B_2 \uplus \dots \uplus B_k, \\ E_{ij} &:= \{v_{\{x,y\}} \in E^* \mid x \in V_i, y \in V_j\}, & \text{and} & W &:= \{w_h \mid 1 \leq h \leq \binom{k}{2} - 1\}. \end{aligned}$$

The edges of G' are as follows. For each edge $e = \{v_i^p, v_j^q\} \in E$, vertex $v_e \in E_{ij}$ of G' is *p*-connected to B_i and *q*-connected to B_j . Vertex $s \in V'$ is adjacent to all vertices in $E_{1,2}$ and

10:10 Parameterized algorithms and data reduction for safe convoy routing

vertex $t \in V'$ is adjacent to all vertices in $E_{k-1,k}$. Finally, to describe the edges incident to vertices in W , consider any ordering of pairs $\{(i, j) \mid 1 \leq i < j \leq k\}$. Then, vertex $w_h \in W$ is adjacent to all vertices in E_{ij} and to all vertices in $E_{i'j'}$, where (i, j) is the h -th pair in the ordering and (i', j') is the $(h+1)$ -st. This finishes the construction.

To prove Theorem 2.14, we show that Construction 2.19 is a polynomial-time many-one reduction that generates bipartite $K_{r,r}$ -free graphs with $r + \text{vc} \in \text{poly}(k \log n)$.

► **Lemma 2.20.** *The graph created by Construction 2.19 from an n -vertex instance $G = (V_1 \uplus V_2 \uplus \dots \uplus V_k, E)$ of MULTICOLORED CLIQUE is bipartite, $K_{r,r}$ -free for $r := 2k \log n + \binom{k}{2} + 2$, and admits a vertex cover of size $r - 1$.*

Proof. The constructed graph $G' = (V', E')$ is bipartite with $V' = X \uplus Y$, where

$$X = \{s, t\} \cup W \cup B \quad \text{and} \quad Y = N(B) \cup E^*.$$

Hence, X is a vertex cover of size at most $r - 1$ in G' . Finally, consider any $K_{r,r}$ whose vertex set is partitioned into two independent sets $X' \uplus Y' \subseteq V'$. Since $|X'| = |Y'| = r$, $|X' \cap X| \leq r - 1$, and $|Y' \cap X| \leq r - 1$, we find $u \in X' \cap Y$ and $v \in Y' \cap Y$. Observe that $\{u, v\}$ is an edge in the $K_{r,r}$ but not in G' . Thus, the $K_{r,r}$ is not a subgraph of G' . ◀

► **Lemma 2.21.** *Construction 2.19 is a polynomial parameter transformation of MULTICOLORED CLIQUE parameterized by $k \log n$ into SSP in $K_{r,r}$ -free graphs parameterized by $\text{vc} + r$.*

Proof. Let $I' := (G', s, t, k', \ell')$ be the SSP instance created by Construction 2.19 from an MULTICOLORED CLIQUE instance $G = (V, E)$. In Lemma 2.20, we already showed $\text{vc} + r \in \text{poly}(k \log n)$. Thus, it remains to show that G is a yes-instance if and only if I' is.

(\Rightarrow) Let C be the edge set of a clique of size k in G . For each $1 \leq i < j \leq k$, C contains exactly one edge e between V_i and V_j . Thus, $E_C := \{v_e \in E^* \mid e \in C\}$ is a set of $\binom{k}{2}$ vertices—exactly one vertex of E_{ij} for each $1 \leq i < j \leq k$. Thus, by Construction 2.19, G' contains an s - t -path $P = (V_P, E_P)$ with $|V_P| \leq k'$: its inner vertices are $E_C \cup W$, alternating between these two sets. To show that I' is a yes-instance, it remains to show $|N(V_P)| \leq \ell'$.

Since P contains all vertices of W , one has $N(V_P) \subseteq B \cup (E^* \setminus E_C)$, where $|E^* \setminus E_C| = |E| - \binom{k}{2}$. To show $|N(V_P)| \leq \ell'$, it remains to show that $|N(V_P) \cap B| \leq k \log \tilde{n}$. To this end, we show that $|N(V_P) \cap B_i| \leq \log \tilde{n}$ for each $i \in \{1, \dots, k\}$.

The vertices in $W \cup \{s, t\}$ have no neighbors in B . Thus, consider arbitrary vertices $v_{e_1}, v_{e_2} \in E_C$ such that $N(v_{e_1}) \cap B_i \neq \emptyset$ and $N(v_{e_2}) \cap B_i \neq \emptyset$ for some $i \in \{1, \dots, k\}$ (possibly, $e_1 = e_2$). Then, $e_1 = \{v_i^p, v_j^q\}$ and $e_2 = \{v_i^{p'}, v_j^{q'}\}$. Since C is a clique, e_1 and e_2 are incident to the same vertex of V_i . Thus, we have $p = p'$. Both v_{e_1} and v_{e_2} are therefore p -connected to B_i and hence have the same $\log \tilde{n}$ neighbors in B_i . It follows that $|N(V_P) \cap B_i| \leq \log \tilde{n}$ and, consequently, that I' is a yes-instance.

(\Leftarrow) Let $P = (V_P, E_P)$ be an s - t -path in G' with $|V_P| \leq k'$ and $|N(V_P)| \leq \ell'$. The path P does not contain any vertex of B , since each of them has $\ell' + 1$ neighbors of degree one. Thus, the inner vertices of P alternate between vertices in W and in E^* and we get $N(V_P) = (E^* \setminus V_P) \cup (N(V_P) \cap B)$. Since P contains one vertex of E_{ij} for each $1 \leq i < j \leq k$, we know $|E^* \setminus V_P| = |E| - \binom{k}{2}$. Thus, since $|N(V_P)| \leq \ell'$, we have $|N(V_P) \cap B| \leq k \log \tilde{n}$. We exploit this to show that the set $C := \{e \in E \mid v_e \in V_P \cap E^*\}$ is the edge set of a clique in G . To this end, it is enough to show that, for each $i \in \{1, \dots, k\}$, any two edges $e_1, e_2 \in C$ with $e_1 \cap V_i \neq \emptyset$ and $e_2 \cap V_i \neq \emptyset$ have the same endpoint in V_i : then C is a set of $\binom{k}{2}$ edges on k vertices and thus forms a k -clique.

For each $1 \leq i < j \leq k$, P contains exactly one vertex $v \in E_{ij}$, which has exactly $\log \tilde{n}$ neighbors in each of B_i and B_j . Thus, from $|N(V_P) \cap B| \leq k \log \tilde{n}$ follows $|N(V_P) \cap B_i| = \log \tilde{n}$ for each $i \in \{1, \dots, k\}$. It follows that, if two vertices v_{e_1} and v_{e_2} on P both have neighbors in B_i , then both are p -connected to B_i for some p , which means that the edges e_1 and e_2 of G share endpoint v_i^p .

We conclude that C is the edge set of a clique of size k in G . Hence, G is a yes-instance. ◀

To prove Theorem 2.14, it is now a matter of putting together Lemma 2.21 and the fact that MULTICOLORED CLIQUE parameterized by $k \log n$ is WK[1]-complete.

Proof of Theorem 2.14. By Lemma 2.21, Construction 2.19 is a polynomial parameter transformation from MULTICOLORED CLIQUE parameterized by $k \log n$ to SSP parameterized by $vc + r$ in $K_{r,r}$ -free graphs.

MULTICOLORED CLIQUE parameterized by $k \log n$ is known to be WK[1]-complete [31] and hence, does not admit a polynomial-size problem kernel unless $\text{coNP} \subseteq \text{NP/poly}$. From the polynomial parameter transformation in Construction 2.19, it thus follows that SSP is WK[1]-hard parameterized by $vc + r$ and does not admit a polynomial-size problem kernel unless $\text{coNP} \subseteq \text{NP/poly}$. ◀

3 Tree-like graphs

In this section, we present results for SSP in tree-like graphs. Such graphs naturally arise as waterways: when ignoring the few man-made canals, the remaining, natural waterways usually form a forest [23].

Moreover, graphs of small *treewidth* (formally defined in Section 3.1) are interesting since, as described in Section 2.1, graphs with constant crossing number have treewidth at most \sqrt{q} for many graph parameters q . Thus, one can derive subexponential-time algorithms for these parameters from single-exponential algorithms for treewidth, like we did in Section 2.1.

First, in Section 3.1, we describe an algorithm that efficiently solves SSP on graphs of small treewidth. Second, in Section 3.2, we show that SSP allows for no problem kernel with size polynomial in the treewidth of the input graph. Third, in Section 3.3, we complement this negative result by a problem kernel with size polynomial in the feedback edge number of the input graph.

3.1 Fixed-parameter algorithm for graphs with small treewidth

In this section, we sketch a $2^{O(\text{tw})} \cdot \ell^2 \cdot n$ -time algorithm for SSP in graphs of treewidth tw , which will also conclude the proof of the $2^{O(\sqrt{n})}$ -time algorithm for SSP in graphs with constant crossing number (Theorem 2.1). Before describing the algorithm, we formally introduce the treewidth concept.

► **Definition 3.1** (tree decomposition, treewidth). A *tree decomposition* $\mathbb{T} = (T, \beta)$ of a graph $G = (V, E)$ consists of a tree T and a function $\beta: V(T) \rightarrow 2^V$ that associates each *node* x of the tree T with a subset $B_x := \beta(x) \subseteq V$, called a *bag*, such that

- i) for each vertex $v \in V$, there is a node x of T with $v \in B_x$,
- ii) for each edge $\{u, v\} \in E$, there is a node x of T with $\{u, v\} \subseteq B_x$,
- iii) for each $v \in V$ the nodes x with $v \in B_x$ induce a subtree of T .

The *width* of \mathbb{T} is $w(\mathbb{T}) := \max_{x \in V(T)} |B_x| - 1$. The *treewidth* of G is $\text{tw}(G) := \min\{w(\mathbb{T}) \mid \mathbb{T} \text{ is a tree decomposition of } G\}$.

► **Theorem 3.2.** SHORT SECLUDED PATH is solvable in $2^{O(tw)} \cdot \ell^2 \cdot n$ time in graphs of treewidth tw .

Bodlaender et al. [9] proved that a tree decomposition of width $O(tw(G))$ of a graph G is computable in $2^{O(tw)} \cdot n$ -time. Applying the following Proposition 3.3 to such a tree decomposition yields Theorem 3.2:

► **Proposition 3.3** (★). VERTEX-WEIGHTED SHORT SECLUDED PATH is solvable in $n \cdot \ell^2 \cdot tw^{O(1)} \cdot (2 + 12 \cdot 2^\omega)^{tw}$ time when a tree decomposition of width tw is given, where $\omega < 2.2373$ is the matrix multiplication exponent.

To prove Theorem 3.2, it thus remains to prove Proposition 3.3. Note that Proposition 3.3 actually solves the weighted problem VW-SSP (Problem 2.6), where the term ℓ^2 is only pseudo-polynomial for VW-SSP. It is a true polynomial for SSP since we can assume $\ell \leq n$.

3.1.1 Assumptions on the tree decomposition

Our algorithm will work on simplified tree decompositions, which can be obtained from a classical tree decomposition of width tw in $n \cdot tw^{O(1)}$ time without increasing its width [8].

► **Definition 3.4** (nice tree decomposition). A nice tree decomposition \mathbb{T} is a tree decomposition with one special bag r called *the root* and in which each bag is of one of the following types.

Leaf node: a leaf x of \mathbb{T} with $B_x = \emptyset$.

Introduce vertex node: an internal node x of \mathbb{T} with one child y such that $B_x = B_y \cup \{v\}$ for some vertex $v \notin B_y$. This node is said to *introduce vertex* v .

Introduce edge node: an internal node x of \mathbb{T} labeled with an edge $\{u, v\} \in E$ and with one child y such that $\{u, v\} \subseteq B_x = B_y$. This node is said to *introduce edge* $\{u, v\}$.

Forget node: an internal node x of \mathbb{T} with one child y such that $B_x = B_y \setminus \{v\}$ for some node $v \in B_y$. This node is said to *forget* v .

Join node: an internal node x of \mathbb{T} with two children y and z such that $B_x = B_y = B_z$.

We additionally require that each edge is introduced at most once and make the following, problem specific assumptions on tree decompositions.

► **Assumption 3.5.** When solving VW-SSP, we will assume that the source s and destination t of the sought path are contained in all bags of the tree decomposition and that the root bag contains only s and t . This ensures that

- every bag contains vertices of the sought solution, and that
- s and t are never forgotten nor introduced.

Such a tree decomposition can be obtained from a nice tree decomposition by rooting it at a leaf (an empty bag) and adding s and t to all bags. This will increase the width of the tree decomposition by at most two.

Our algorithm will be based on computing partial solutions for subgraphs induced by a node of a tree decomposition by means of combining partial solutions for the subgraphs induced by its children. Formally, these subgraphs are the following.


► **Definition 3.6** (subgraphs induced by a tree decomposition). Let $G = (V, E)$ be a graph and \mathbb{T} be a nice tree decomposition for G with root r . Then, for any node x of \mathbb{T} ,

$$V_x := \{v \in V \mid v \in B_y \text{ for a descendant } y \text{ of } x\}, \text{ and}$$

$$G_x := (V_x, E_x), \text{ where } E_x = \{e \in E \mid e \text{ is introduced in a descendant of } x\}.$$

Herein, we consider each node x of \mathbb{T} to be a descendent of itself.

Having defined subgraphs induced by subtrees, we can define partial solutions in them.



■ **Figure 3.1** Illustration of a partial solution: the (blue) thick edges are an overall solution, where the darker edges are the part of the solution in G_x . The (red) dashed edges are forbidden to exist.

3.1.2 Partial solutions

Assume that we have a solution path P to VW-SSP. Then, the part of P in G_x is a collection \mathcal{P} of paths (some might consist of a single vertex). When computing a partial solution for a parent y of x , we ideally want to check which partial solutions for x can be continued to partial solutions for y . However, we cannot try all possible partial solutions for G_x —there might be too many. Moreover, this is not necessary: by Definition 3.1(ii)–(iii), vertices in bag B_y cannot be vertices of and cannot have edges to vertices of $V_x \setminus B_x$. Thus, it is enough to know the states of vertices in bag B_x in order to know which partial solutions of x can be continued to y . The state of such vertices is characterized by

- which vertices of B_x are end points of paths in \mathcal{P} , inner vertices of paths in \mathcal{P} , or paths of zero length in \mathcal{P} ,
- which vertices of B_x are allowed to be neighbors of the solution path P ,
- how many neighbors the solution path P is allowed to have in G_x , and
- which vertices of B_x belong to the same path of \mathcal{P} .

► **Definition 3.7** (partial solution). Let $(G, s, t, k, \ell, \kappa, \lambda)$ be an instance of VW-SSP. For a set \mathcal{P} of paths in G and a set N of vertices in G , let

$$\Lambda(\mathcal{P}, N) := \sum_{P \in \mathcal{P}} \sum_{v \in N(V(P))} \lambda(v) + \sum_{v \in N} \lambda(v) \quad \text{and} \quad K(\mathcal{P}) := \sum_{P \in \mathcal{P}} \sum_{v \in V(P)} \kappa(v).$$

Moreover, let \mathbb{T} be a tree decomposition for G , x be a node of \mathbb{T} , $D_z \uplus D_e \uplus D_i \uplus N \subseteq B_x$ such that $\{s, t\} \subseteq D_z \cup D_e$, p be a partition of $D := D_z \cup D_e \cup D_i$, and $l \leq \ell$.

Then, we call (D_z, D_e, D_i, N, l) a *pre-signature* and $S = (D_z, D_e, D_i, N, l, p)$ a *solution signature* at x . A set \mathcal{P} of paths in G_x is a *partial solution* of cost $K(\mathcal{P})$ for S if

- i) D_z are exactly the vertices of zero-length paths $P \in \mathcal{P}$,
- ii) D_e are exactly the end points of non-zero-length paths $P \in \mathcal{P}$,
- iii) D_i are exactly those vertices in B_x that are inner vertices of paths $P \in \mathcal{P}$,
- iv) for each path $P \in \mathcal{P}$, $N(V(P)) \cap B_x \subseteq N$,
- v) $\Lambda(\mathcal{P}, N) \leq l$, and
- vi) \mathcal{P} consists of exactly $|p|$ paths such that each two vertices $u, v \in D$ belong to the same path of \mathcal{P} if and only if they are in the same set of the partition p .

For a solution signature S at a node x , we denote

$$\mathcal{E}_x(S) := \{\mathcal{P} \mid \mathcal{P} \text{ is a partial solution for } S\},$$

$$\min K_x(S) := \min\{K(\mathcal{P}) \mid \mathcal{P} \in \mathcal{E}_x(S)\}.$$

10:14 Parameterized algorithms and data reduction for safe convoy routing

Because of Assumption 3.5, our input instance to VW-SSP is a yes-instance if and only if

$$\min K_r(\emptyset, \{s, t\}, \emptyset, \emptyset, \ell, \{\{s, t\}\}) \leq k. \quad (3.1)$$

Therefore, our aim is computing this cost. The naive dynamic programming approach is:

- compute $\min K_x(S)$ for each solution signature S and each leaf node x ,
- compute $\min K_x(S)$ for each solution signature S and each inner node x under the assumptions that $\min K_y(S')$ has already been computed for all solution signatures S' at children y of x .

However, this approach is not suitable to prove Proposition 3.3, since the number of possible solution signatures is too large: the number of different partitions p of tw vertices is the tw -th Bell number, whose best known upper bound is $O(\text{tw}^{\text{tw}}/\log \text{tw})$.

3.1.3 Reducing the number of partitions

To reduce the number of needed partitions, we use an approach developed by Bodlaender et al. [8], which also proved its effectivity in experiments [18]. We will replace the task of computing (3.1) for all possible partitions by computing only sets of *weighted partitions* containing the needed information.

► **Definition 3.8** (sets of weighted partitions). Let $\Pi(U)$ be the set of all partitions of U . A *set of weighted partitions* is a set $\mathcal{A} \subseteq \Pi(U) \times \mathbb{N}$. For a *weighted partition* $(p, w) \in \mathcal{A}$, we call w its *weight*.

Using sets of weighted partitions, we can reformulate our task of computing $\min K_x(S)$ for all bags B_x and all solution signatures S as follows. Consider a pre-signature $S = (D_z, D_i, D_e, N, \ell)$ for a node x of a tree decomposition. Then, for each $p \in \Pi(D_z \cup D_i \cup D_e)$, (S, p) is a solution signature. Thus, we can consider

$$\mathcal{A}_x(S) := \left\{ \left(p, \min_{\mathcal{P} \in \mathcal{E}_x(S, p)} K(\mathcal{P}) \right) \mid p \in \Pi(D_z \cup D_i \cup D_e) \wedge \mathcal{E}_x(S, p) \neq \emptyset \right\}. \quad (3.2)$$

Now, our problem of verifying (3.1) at the root node r of a tree decomposition is equivalent to checking whether $\mathcal{A}_r(\emptyset, \{s, t\}, \emptyset, \emptyset, \ell)$ contains a partition $\{\{s, t\}\}$ of weight at most k . Thus we can, in a classical dynamic programming manner

- compute $\mathcal{A}_x(S)$ for each pre-signature S and each leaf node x ,
- compute $\mathcal{A}_x(S)$ for each pre-signature S and each inner node x under the assumption that $\mathcal{A}_y(S')$ has already been computed for all pre-signatures S' at children y of x .

Yet we will not work with the full sets $\mathcal{A}_x(S)$ but with “representative” subsets of size $2^{O(\text{tw})}$. Since the number of pre-signatures is $2^{O(\text{tw})} \cdot \ell$, this will allow us to prove Proposition 3.3.

In order to describe the intuition behind representative sets of weighted partitions, we need some notation.

► **Definition 3.9** (partition lattice). The set $\Pi(U)$ is semi-ordered by the coarsening relation \sqsubseteq , where $p \sqsubseteq q$ if every set of p is included in some set of q . We also say that q is *coarser* than p and that p is *finer* than q .

For two partitions $p, q \in \Pi(U)$, by $p \sqcup q$ we denote the (unique) finest partition that is coarser than both p and q .

To get an intuition for the $p \sqcup q$ operation, recall from Definition 3.7 that we will use a partition p to represent connected components of partial solutions: two vertices are connected if and only if they are in the same set of p . In these terms, if $p \in \Pi(U)$ are the vertex sets of the connected components of a graph (U, E) and $q \in \Pi(U)$ are the vertex sets of the connected components of a graph (U, E') , then $p \sqcup q$ are the vertex sets of the connected components of the graph $(U, E \cup E')$.

Now, assume that there is a solution P to VW-SSP in a graph G and consider an arbitrary node x of a tree decomposition. Then, the subpaths \mathcal{P} of P that lie in G_x are a partial solution for some solution signature (D_z, D_e, D_i, N, l, p) at x . The partition p of $D := D_z \cup D_e \cup D_i$ consists of the sets of vertices of D that are connected by paths in \mathcal{P} . Since, in the overall solution P , the vertices in D are all connected, the vertices of D are connected in $G \setminus E_x$ according to a partition q of D such that $p \sqcup q = \{D\}$. Now, if in P , we replace the subpaths \mathcal{P} by any other partial solution \mathcal{P}' to a solution signature $(D_z, D_e, D_i, N, l, p')$ such that $K(\mathcal{P}') \leq K(\mathcal{P})$ and $p' \sqcup q = \{D\}$, then we obtain a solution P' for G with at most the cost of P . Thus, one of the two weighted partitions $(p, K(p))$ and $(p', K(p'))$ in $\mathcal{A}_x(D_z, D_e, D_i, N, l)$ is redundant.

This concept of redundancy can be formalized as *representative sets* and representative sets of size $2^{O(\text{tw})}$ can be efficiently computed using results of Bodlaender et al. [8]. To prove Proposition 3.3, it is enough to derive a recurrence relation for (3.2) that plays well together with the framework of Bodlaender et al. [8].

3.2 Hardness of kernelization for graphs of small treewidth

In the previous section, we have seen that SSP is efficiently solvable in tree-like graphs, namely, in graphs of small treewidth. We can complement this result as follows.

► **Theorem 3.10** (\star). *SHORT SECLUDED PATH has no problem kernel with size polynomial in $\text{tw} + k + \ell$, even on planar graphs with maximum degree six, where tw is the treewidth, unless $\text{coNP} \subseteq \text{NP/poly}$ and the polynomial-time hierarchy collapses to the third level.*

To prove Theorem 3.10, we use a special kind of reduction called *cross composition* [7].

► **Definition 3.11** (cross composition). *A polynomial equivalence relation \sim is an equivalence relation over Σ^* such that*

- there is an algorithm that decides $x \sim y$ in polynomial time for any two instances $x, y \in \Sigma^*$, and such that
- the number of equivalence classes of \sim over any *finite* set $S \subseteq \Sigma^*$ is polynomial in $\max_{x \in S} |x|$.

A language $K \subseteq \Sigma^*$ *cross-composes* into a parameterized language $L \subseteq \Sigma^* \times \mathbb{N}$ if there is a polynomial-time algorithm, called *cross composition*, that, given a sequence x_1, \dots, x_p of p instances that are equivalent under some polynomial equivalence relation, outputs an instance (x^*, k) such that

- k is bounded by a polynomial in $\max_{i=1}^p |x_i| + \log p$ and
- $(x^*, k) \in L$ if and only if there is an $i \in \{1, \dots, p\}$ such that $x_i \in K$.

Cross compositions can be used to rule out problem kernels of polynomial size using the following result of Bodlaender et al. [7].

► **Proposition 3.12** (Bodlaender et al. [7]). *If a NP-hard language $K \subseteq \Sigma^*$ cross-composes into the parameterized language $L \subseteq \Sigma^* \times \mathbb{N}$, then there is no polynomial-size problem kernel for L unless $\text{coNP} \subseteq \text{NP/poly}$ and the polynomial-time hierarchy collapses to the third level.*

Using a cross composition, Luckow and Fluschnik [37] proved that SSP on planar graphs of maximum degree six does not admit a problem kernel with size polynomial in $k + \ell$. To prove Theorem 3.10, one can show that the graph created by their cross composition has treewidth at most $3n + 3$, where n is the number of vertices in each input instance to their cross composition.

3.3 Effective data reduction for graphs with small feedback edge set

In the previous section, we have seen that SSP has no problem kernel with size polynomial in the treewidth of the input graph. We can complement this result by proving a polynomial-size problem kernel for another parameter that measures the tree-likeness of a graph: the *feedback edge number* of a graph is the smallest number of edges one has to delete to obtain a forest. Formally, we can prove the following theorem.

► **Theorem 3.13** (\star). *SHORT SECLUDED PATH has a problem kernel with size polynomial in the feedback edge number of the input graph.*

The outline of the proof of Theorem 3.13 is similar that of the proof of Theorem 2.5: we first, in linear time, produce a weighted instance with $O(\text{fes})$ vertices, then reduce the weights using Lemma 2.13, and finally transform the weighted instance back into an instance for SSP.

Towards the first step, we apply data reduction rules that reduce the number of degree-one vertices and the length of paths of degree-two vertices to $O(\text{fes})$. Because of Reduction Rule 1.4, the graph without the fes edges of a feedback edge set is a tree. Thus, its overall number of vertices and edges will be bounded by $O(\text{fes})$.

4 Conclusion

Concluding, we point out that our algorithms for VW-SSP on graphs of bounded treewidth (Theorem 3.2) can easily be generalized to a problem variant where also edges have a weight counting towards the path length, and so can our subexponential-time algorithms in planar graphs (Theorem 2.1). Moreover, the technique of Bodlaender et al. [8] that our algorithm is based on has experimentally been proven to be practically implementable [18].

In contrast, we observed SSP to be a problem for which provably effective polynomial-time data reduction is rather hard to obtain (Theorems 2.14 and 3.10). Therefore, studying relaxed models of data reduction with performance guarantees like approximate [19, 36] or randomized kernelization [35] seems worthwhile.

Indeed, our few positive results on kernelization, that is, our problem kernels of size $\text{vc}^{O(r)}$ in $K_{r,r}$ -free graphs and of size $\text{fes}^{O(1)}$ in graphs of feedback edge number fes for SSP (Theorems 2.5 and 3.13), for now, can be mainly seen as a proof of concept, since they employ the quite expensive weight reduction algorithm of Frank and Tardos [22] and we have no “direct” way of reducing VW-SSP back to SSP. On the positive side, our solution algorithms also work for VW-SSP, so that reducing weights or reducing back to SSP may be unnecessary from a practical point of view.

References

- 1 Hannah Bast, Daniel Delling, Andrew Goldberg, Matthias Müller-Hannemann, Thomas Pajor, Peter Sanders, Dorothea Wagner, and Renato F. Werneck. Route planning in transportation networks. In *Algorithm Engineering: Selected Results and Surveys*, volume 9220 of *Lecture Notes in Computer Science*, pages 19–80. Springer, 2016. [10.1007/978-3-319-49487-6_2](https://doi.org/10.1007/978-3-319-49487-6_2).
- 2 René van Bevern, Rolf Niedermeier, Manuel Sorge, and Mathias Weller. Complexity of arc routing problems. In *Arc Routing: Problems, Methods, and Applications*, volume 20 of *MOS-SIAM Series on Optimization*. SIAM, 2014. [10.1137/1.9781611973679.ch2](https://doi.org/10.1137/1.9781611973679.ch2).
- 3 René van Bevern, Christian Komusiewicz, and Manuel Sorge. A parameterized approximation algorithm for the mixed and windy capacitated arc routing problem: Theory and experiments. *Networks*, 70(3):262–278, 2017. [10.1002/net.21742](https://doi.org/10.1002/net.21742). WARP 2 special issue.
- 4 René van Bevern, Till Fluschnik, George B. Mertzios, Hendrik Molter, Manuel Sorge, and Ondřej Suchý. The parameterized complexity of finding secluded solutions to some classical optimization problems on graphs. *Discrete Optimization*, 2018. [10.1016/j.disopt.2018.05.002](https://doi.org/10.1016/j.disopt.2018.05.002). In press, available on arXiv:1606.09000v5.
- 5 Hans-Joachim Böckenhauer, Juraj Hromkovič, Joachim Kneis, and Joachim Kupke. The parameterized approximability of TSP with deadlines. *Theory of Computing Systems*, 41(3):431–444, 2007. [10.1007/s00224-007-1347-x](https://doi.org/10.1007/s00224-007-1347-x).
- 6 Hans L. Bodlaender, Rodney G. Downey, Michael R. Fellows, and Danny Hermelin. On problems without polynomial kernels. *Journal of Computer and System Sciences*, 75(8):423–434, 2009. [10.1016/j.jcss.2009.04.001](https://doi.org/10.1016/j.jcss.2009.04.001).
- 7 Hans L. Bodlaender, Bart M. P. Jansen, and Stefan Kratsch. Kernelization lower bounds by cross-composition. *SIAM Journal on Discrete Mathematics*, 28(1):277–305, 2014. [10.1137/120880240](https://doi.org/10.1137/120880240).
- 8 Hans L. Bodlaender, Marek Cygan, Stefan Kratsch, and Jesper Nederlof. Deterministic single exponential time algorithms for connectivity problems parameterized by treewidth. *Information and Computation*, 243:86–111, 2015. [10.1016/j.ic.2014.12.008](https://doi.org/10.1016/j.ic.2014.12.008).
- 9 Hans L. Bodlaender, Pål Grønås Drange, Markus S. Dregi, Fedor V. Fomin, Daniel Lokshtanov, and Michał Pilipczuk. A $c^k n$ 5-approximation algorithm for treewidth. *SIAM Journal on Computing*, 45(2):317–378, 2016. [10.1137/130947374](https://doi.org/10.1137/130947374).
- 10 Drago Bokal, Gasper Fijavz, and Bojan Mohar. The minor crossing number. *SIAM Journal on Discrete Mathematics*, 20(2):344–356, 2006. [10.1137/05062706X](https://doi.org/10.1137/05062706X).
- 11 Shiri Chechik, Matthew P. Johnson, Merav Parter, and David Peleg. Secluded connectivity problems. *Algorithmica*, 79(3):708–741, 2017. [10.1007/s00453-016-0222-z](https://doi.org/10.1007/s00453-016-0222-z).
- 12 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015. [10.1007/978-3-319-21275-3](https://doi.org/10.1007/978-3-319-21275-3).
- 13 Erik D. Demaine and Mohammadtaghi Hajiaghayi. Linearity of grid minors in treewidth with applications through bidimensionality. *Combinatorica*, 28(1):19–36, 2008. [10.1007/s00493-008-2140-4](https://doi.org/10.1007/s00493-008-2140-4).
- 14 Reinhard Diestel. *Graph Theory*, volume 173 of *Graduate Texts in Mathematics*. Springer, 4th edition, 2010. [10.1007/978-3-662-53622-3](https://doi.org/10.1007/978-3-662-53622-3).
- 15 Frederic Dorn, Hannes Moser, Rolf Niedermeier, and Mathias Weller. Efficient algorithms for Eulerian Extension and Rural Postman. *SIAM Journal on Discrete Mathematics*, 27(1):75–94, 2013. [10.1137/110834810](https://doi.org/10.1137/110834810).
- 16 Rod G. Downey and Michael R. Fellows. *Fundamentals of Parameterized Complexity*. Springer, 2013. [10.1007/978-1-4471-5559-1](https://doi.org/10.1007/978-1-4471-5559-1).

- 17 Michael Etscheid, Stefan Kratsch, Matthias Mnich, and Heiko Röglin. Polynomial kernels for weighted problems. *Journal of Computer and System Sciences*, 84(Supplement C): 1–10, 2017. [10.1016/j.jcss.2016.06.004](https://doi.org/10.1016/j.jcss.2016.06.004).
- 18 Stefan Fafanie, Hans L. Bodlaender, and Jesper Nederlof. Speeding up dynamic programming with representative sets: An experimental evaluation of algorithms for steiner tree on tree decompositions. *Algorithmica*, 71(3):636–660, 2015. [10.1007/s00453-014-9934-0](https://doi.org/10.1007/s00453-014-9934-0).
- 19 Michael R. Fellows, Ariel Kulik, Frances A. Rosamond, and Hadas Shachnai. Parameterized approximation via fidelity preserving transformations. *Journal of Computer and System Sciences*, 93:30–40, 2018. [10.1016/j.jcss.2017.11.001](https://doi.org/10.1016/j.jcss.2017.11.001).
- 20 Jörg Flum and Martin Grohe. *Parameterized Complexity Theory*. Springer, 2006. [10.1007/3-540-29953-X](https://doi.org/10.1007/3-540-29953-X).
- 21 Fedor V. Fomin, Petr A. Golovach, Nikolay Karpov, and Alexander S. Kulikov. Parameterized complexity of secluded connectivity problems. *Theory of Computing Systems*, 61(3):795–819, 2017. [10.1007/s00224-016-9717-x](https://doi.org/10.1007/s00224-016-9717-x).
- 22 András Frank and Éva Tardos. An application of simultaneous diophantine approximation in combinatorial optimization. *Combinatorica*, 7(1):49–65, 1987. [10.1007/BF02579200](https://doi.org/10.1007/BF02579200).
- 23 Achille Giacometti. River networks. In *Complex Networks*, Encyclopedia of Life Support Systems (EOLSS), pages 155–180. EOLSS Publishers/UNESCO, 2010.
- 24 Petr A. Golovach, Pinar Heggernes, Paloma T. Lima, and Pedro Montealegre. Finding connected secluded subgraphs. In *Proceedings of the 12th International Symposium on Parameterized and Exact Computation, IPEC 2017, September 6-8, 2017, Vienna, Austria*, volume 89 of *LIPICs*, pages 18:1–18:13. Schloss Dagstuhl – Leibniz-Zentrum fuer Informatik, 2017. [10.4230/LIPICs.IPEC.2017.18](https://doi.org/10.4230/LIPICs.IPEC.2017.18).
- 25 Jiong Guo and Rolf Niedermeier. Invitation to data reduction and problem kernelization. *ACM SIGACT News*, 38(1):31–45, 2007. [10.1145/1233481.1233493](https://doi.org/10.1145/1233481.1233493).
- 26 G. Gutin and V. Patel. Parameterized traveling salesman problem: Beating the average. *SIAM Journal on Discrete Mathematics*, 30(1):220–238, 2016. [10.1137/140980946](https://doi.org/10.1137/140980946).
- 27 Gregory Gutin, Gabriele Muciaccia, and Anders Yeo. Parameterized complexity of k -Chinese Postman Problem. *Theoretical Computer Science*, 513:124–128, 2013. [10.1016/j.tcs.2013.10.012](https://doi.org/10.1016/j.tcs.2013.10.012).
- 28 Gregory Gutin, Mark Jones, and Magnus Wahlström. The mixed chinese postman problem parameterized by pathwidth and treedepth. *SIAM Journal on Discrete Mathematics*, 30(4):2177–2205, 2016. [10.1137/15M1034337](https://doi.org/10.1137/15M1034337).
- 29 Gregory Gutin, Mark Jones, and Bin Sheng. Parameterized complexity of the k -arc chinese postman problem. *Journal of Computer and System Sciences*, 84:107–119, 2017. [10.1016/j.jcss.2016.07.006](https://doi.org/10.1016/j.jcss.2016.07.006).
- 30 Gregory Gutin, Magnus Wahlström, and Anders Yeo. Rural Postman parameterized by the number of components of required edges. *Journal of Computer and System Sciences*, 83(1):121–131, 2017. [10.1016/j.jcss.2016.06.001](https://doi.org/10.1016/j.jcss.2016.06.001).
- 31 Danny Hermelin, Stefan Kratsch, Karolina Sołtys, Magnus Wahlström, and Xi Wu. A completeness theory for polynomial (turing) kernelization. *Algorithmica*, 71(3):702–730, 2015. [10.1007/s00453-014-9910-8](https://doi.org/10.1007/s00453-014-9910-8).
- 32 Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *Journal of Computer and System Sciences*, 63(4): 512–530, 2001. [10.1006/jcss.2001.1774](https://doi.org/10.1006/jcss.2001.1774).
- 33 Philip N. Klein and Daniel Marx. A subexponential parameterized algorithm for Subset TSP on planar graphs. In *Proceedings of the 25th Annual ACM-SIAM Symposium on*

- Discrete Algorithms (SODA'14)*, pages 1812–1830. Society for Industrial and Applied Mathematics, 2014. [10.1137/1.9781611973402.131](https://doi.org/10.1137/1.9781611973402.131).
- 34 Stefan Kratsch. Recent developments in kernelization: A survey. *Bulletin of the EATCS*, 113, 2014. <http://eatcs.org/beatcs/index.php/beatcs/article/view/285>.
- 35 Stefan Kratsch and Magnus Wahlström. Compression via matroids: A randomized polynomial kernel for odd cycle transversal. *ACM Transactions on Algorithms*, 10(4): 20:1–20:15, 2014. [10.1145/2635810](https://doi.org/10.1145/2635810).
- 36 Daniel Lokshtanov, Fahad Panolan, M. S. Ramanujan, and Saket Saurabh. Lossy kernelization. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing (STOC 2017)*, pages 224–237. ACM, 2017. [10.1145/3055399.3055456](https://doi.org/10.1145/3055399.3055456).
- 37 Max-Jonathan Luckow and Till Fluschnik. On the computational complexity of length- and neighborhood-constrained path problems. Available on arXiv:1808.02359, 2018.
- 38 Neeldhara Misra, Venkatesh Raman, and Saket Saurabh. Lower bounds on kernelization. *Discrete Optimization*, 8(1):110–128, 2011. [10.1016/j.disopt.2010.10.001](https://doi.org/10.1016/j.disopt.2010.10.001).
- 39 Rolf Niedermeier. *Invitation to Fixed-Parameter Algorithms*. Oxford University Press, 2006. [10.1093/acprof:oso/9780198566076.001.0001](https://doi.org/10.1093/acprof:oso/9780198566076.001.0001).
- 40 Janos Pach, Rados Radoicic, Gabor Tardos, and Geza Toth. Improving the crossing lemma by finding more crossings in sparse graphs. *Discrete & Computational Geometry*, 36(4):527–552, 2006. [10.1007/s00454-006-1264-9](https://doi.org/10.1007/s00454-006-1264-9).
- 41 Manuel Sorge, René van Bevern, Rolf Niedermeier, and Mathias Weller. From few components to an Eulerian graph by adding arcs. In *Proceedings of the 37th International Workshop on Graph-Theoretic Concepts in Computer Science (WG'11)*, volume 6986 of *Lecture Notes in Computer Science*, pages 307–318. Springer, 2011. [10.1007/978-3-642-25870-1_28](https://doi.org/10.1007/978-3-642-25870-1_28).
- 42 Manuel Sorge, René van Bevern, Rolf Niedermeier, and Mathias Weller. A new view on Rural Postman based on Eulerian Extension and Matching. *Journal of Discrete Algorithms*, 16:12–33, 2012. [10.1016/j.jda.2012.04.007](https://doi.org/10.1016/j.jda.2012.04.007).
- 43 K. Wagner. Über eine Eigenschaft der ebenen Komplexe. *Mathematische Annalen*, 114 (1):570–590, 1937. [10.1007/BF01594196](https://doi.org/10.1007/BF01594196).